

# The effect of local weather conditions and nest box location on the reproduction of the Common Kestrel (*Falco tinnunculus*) in the farmland of eastern Poland

Zbigniew Kasprzykowski\*, Mirosław Rzepała & Artur Golawski

Z. Kasprzykowski, Siedlce University of Natural Sciences and Humanities, Faculty of Exact and Natural Sciences, Prusa 14, 08-110 Siedlce, Poland \* Corresponding author's e-mail: [zbigniew.kasprzykowski@uph.edu.pl](mailto:zbigniew.kasprzykowski@uph.edu.pl)

M. Rzepała, "Stork" Nature Society, Radomska 22/32, 02-323 Warszawa, Poland

A. Golawski, Siedlce University of Natural Sciences and Humanities, Faculty of Exact and Natural Sciences, Prusa 14, 08-110 Siedlce, Poland

Received 11 January 2021, accepted 3 August 2021



Artificial nests like nest boxes are commonly used to improve the reproductive output and population status of wild birds and advance our understanding of their evolutionary ecology. To investigate how the location of nest boxes and local weather conditions affect Common Kestrel reproduction (number of fledglings), 319 broods from 173 nest boxes in east-central Poland were examined. Of the six meteorological parameters analysed, only mean temperature and maximum temperature were selected as the best model. Mean daily temperature had a positive effect but maximum temperature a negative one on Common Kestrel reproduction. The first egg laying date was negatively correlated with the number of fledglings. Contrary to the assumption that trees should offer a superior microclimate in the canopy, nest boxes placed on utility poles have the same numbers of fledglings. The area of the main habitat type is not decisive in Common Kestrel productivity either. These findings may be helpful for ongoing conservation programmes and confirm that utility poles should be the preferred form of deployment of nest boxes as they are easier to monitor and maintain.

## 1. Introduction

Many bird species use man-made structures for nesting, their primary advantage being that they often provide nesting sites in areas where natural ones are not available (Lambrechts *et al.* 2012, Mainwaring 2015, Le Roux *et al.* 2016). Artificial nests like nest boxes are commonly used to improve the reproductive output and population

status of bird species with nest-site limited populations (Golawski *et al.* 2003, Klein *et al.* 2007, Smallwood *et al.* 2009, Rodríguez *et al.* 2011, Libois *et al.* 2012, Schaub *et al.* 2016, Carstens *et al.* 2019) and to improve our understanding of the functional and evolutionary ecology of wild, cavity-nesting birds (Lambrechts *et al.* 2012).

Nest boxes offer shelter against extreme weather (Lambrechts *et al.* 2012). However, weather conditions not only impact breeding

performance directly, but also bring forward the timing of breeding (Carrillo & González-Dávila 2010a, Sumasgutner *et al.* 2020). This could have an influence on factors benefitting birds' reproduction, e.g. higher parental provisioning rates and lower ectoparasite abundance in nests (McDonald *et al.* 2004, Martínez-de la Puente *et al.* 2009). The positioning of nest boxes is also important, as the different compass directions in which they face will expose them to different sets of weather conditions (Butler *et al.* 2009). Generally, studies of falcons confirm that reproductive parameters are to a large extent affected by weather (Dawson & Bortolotti 2000, Rodríguez & Bustamante 2003, Carlzon *et al.* 2018). Nevertheless, if conservation programmes are to be effective, the influence of both nest box location and local weather conditions should be analysed (see Valkama & Korpimäki 1999).

For secondary hole-nesting raptors, the probability that a nest box is occupied and the expression of avian life-history traits of the nest box's occupants depend on aspects of nest box design, location and maintenance procedures (Lambrechts *et al.* 2012). Artificial structures are frequently used by falcons that do not build their own nests, thus benefitting their reproductive parameters (Bortolotti 1994, Fargallo *et al.* 2001). Common Kestrel *Falco tinnunculus* breeding in nest boxes in an open landscape produced more fledglings than its conspecifics breeding in other sites like holes in buildings, pylons and trees (Valkama & Korpimäki 1999, Fargallo *et al.* 2001, Charter *et al.* 2007a). The preference of Common Kestrels for artificial nests is associated with the lower predation rate by Pine Martens *Martes martes*, Goshawks *Accipiter gentilis* and rodents, reduced nest site competition with Long-eared Owls *Asio otus*, and shorter distances from the nest to foraging areas (Valkama & Korpimäki 1999, Fargallo *et al.* 2001). On the other hand, Starlings *Sturnus vulgaris* in North America have been found to usurp nest boxes during the American Kestrel's egg-laying stage, thereby reducing the nesting success of these raptors (Rohrbaugh & Yahner 1997). Nonetheless, nest boxes seem to be safer than natural sites. Despite the higher intensity of nestling infestations in nest boxes, they are less often predated (Fargallo *et al.* 2001). As a consequence, the installation of nest

boxes has led to local increases in the breeding populations of different falcon species and this form of conservation management is often applied in farmland and urban areas (Bux *et al.* 2008, Smallwood & Collopy 2009, Altwegg *et al.* 2014, Costantini & Dell'Omo 2020).

The aim of this research was to investigate how the deployment of nest boxes and local weather conditions affect Common Kestrel reproduction. We hypothesised that the number of fledglings would depend on the nest box site, the time of egg laying, temperature, precipitation and wind speed. We also anticipated that numbers of fledglings would be higher in nest boxes attached to trees rather than utility poles, as the tree canopy would offer a superior microclimate (Valkama & Korpimäki 1999). On the other hand, nest boxes located on utility poles are more exposed to temperature and rainfall, which could affect the number of fledglings (see Costantini *et al.* 2010a, Kreiderits 2016). Knowledge of the impact of these factors may be useful when implementing conservation measures for the Common Kestrel in farmland (see Lambrechts *et al.* 2012).

## 2. Material and methods

### 2.1. Study area

The research was carried out within a radius of ca 50 km around the town of Siedlce (east-central Poland; 52°12'N, 22°17'E). This region has a preponderance of arable land (46.9%), with meadows and pastureland having a 14.7% coverage, and orchards, mainly apple trees, occupying 2.6% of the land. Woodlands cover 24.8% of the region and built-up areas 5.7%. The populations of the towns (Siedlce, Węgrów, Sokołów Podlaski) range from 12 800 to 78 000. The remaining 5.3% consists of water bodies and wasteland (Statistical Yearbook 2015). The area has a temperate transitional climate (Degirmendzic *et al.* 2004).

### 2.2. Data collection

The study was carried out during ten breeding seasons from 2010 to 2019. During this period, 173 nest boxes were visited and 319 breeding

pairs of Common Kestrel investigated. The birds nested in the same type of nest box (30 cm wide × 40 long × 30 cm high), constructed from 2 cm thick pine boards and half-open to one side. The roof was secured with a metal sheet or roofing felt, and all the walls were coated with an impregnating agent. Nest boxes were hung at the same height on trees and on utility poles, between 6 and 7 m above the ground. Data were gathered from 194 nests (93 boxes) deployed on utility poles and from 125 nests (80 boxes) on trees. The mean number of nests monitored each year was 31.9 ( $SD = 19.46$ ). During the breeding season each nest box was inspected at least twice: once in May and then again on the day of the last inspection, depending on the age of the fledglings and the possibility of counting them, since their number was key to this study. If, during the second inspection, the fledglings were too small to be counted, a third inspection was undertaken. The size of the fledged brood was taken to be the number of nestlings in successful nests in the fourth week (Kreiderits *et al.* 2016). To eliminate the year effect (Costantini *et al.* 2010b), the data were standardised against the first egg date (FED) in the population in each year. For all clutches, the first egg date was established by backdating from the day the chicks left the nest, assuming 30 days for incubation (starting with the 3<sup>rd</sup> egg), and the hatching day was based on a 30-day period for the nestling stage. The dominant habitat (meadows or arable land) within a 1 km radius of the boxes was determined using QGIS software. In the open

areas of Europe, Kestrels are known to forage within a few km of the nest, but the mostly do so no farther away than 1000 m (Korpimäki *et al.* 1996, Norrdahl & Korpimäki 2002). The meteorological data were obtained from the weather station in Siedlce (52°25' N, 22°26' E), which is representative of the whole study area. The mean annual temperature during the study period varied from 15.8 to 19.3 °C. Six meteorological parameters were calculated for the nestling-rearing period: three temperature parameters (mean, maximum and minimum), two precipitation parameters (total and day numbers) and wind speed (Table 1). The mean temperature ( $T_{\text{mean}}$ ) was calculated from the mean daily temperature for all the days in the chick-rearing period. The minimum ( $T_{\text{min}}$ ) and maximum temperatures ( $T_{\text{max}}$ ) were the lowest and highest temperatures recorded in that given period. The total precipitation ( $PP_{\text{sum}}$ ) was the total amount of precipitation that fell in a 30-day period, while the number of days with precipitation ( $PP_{\text{days}}$ ) was the total number of days when rain fell. The mean wind speed ( $W_{\text{speed}}$ ) was the average of the mean daily wind speeds in a given period.

### 2.3. Statistical analyses

A General Linear Mixed Model (GLMM) with identity link function and normal error distribution was used to test the hypothesis stated in the introduction. The number of 4-week old

Table 1. Factors for analysing reproduction of Common Kestrel (*Falco tinnunculus*) breeding in nest boxes in eastern Poland.

Code	Description
FED	Standardised first egg laying date
Location	Category of nest box location: utility pole and tree
Habitat	Category of habitat: meadow and arable land
$T_{\text{mean}}$	mean temperature (°C), range 12.8–21.7
$T_{\text{min}}$	minimum temperature (°C), range (–2.4)–9.4
$T_{\text{max}}$	maximum temperature (°C), range 26.3–33.7
$PP_{\text{sum}}$	total precipitation (mm), range 15.5–223.0
$PP_{\text{days}}$	number of days with precipitation (N), range 6–23
$W_{\text{speed}}$	mean wind speed (km/h), range 6.8–11.1

Table 2. Results of three models describing the influence of habitat factors on the numbers of Common Kestrel *Falco tinnunculus* fledglings in eastern Poland. Degrees of freedom (*df*), model log-likelihood (LL), corrected AIC criterion ( $AIC_c$ ), difference between the model and the best model in the data set ( $\Delta AIC_c$ ) and weight for the model ( $AIC_c wt$ ) are shown.

Model (fixed effects)	<i>df</i>	LL	$AIC_c$	$\Delta AIC_c$	$AIC_c wt$
Intercept+FED+Tmean+Tmax	6	-476.933	966.1	0.00	0.081
Intercept+FED+Tmean+Tmax+Habitat	7	-476.374	967.1	0.97	0.049
Intercept+FED+Tmean+Tmax+PPsum	7	-476.635	967.6	1.50	0.038
Intercept+FED+Tmean+Tmax+Site	7	-476.843	968.0	1.91	0.031
Intercept+FED+Tmean+Tmax+Tmin	7	-476.872	968.1	1.97	0.030

nestlings was treated as a dependent variable. Six weather parameters, the first egg laying date, habitat category and nest box location category were independent variables. Nest box identity was introduced as a random factor to account for within-subject consistency in the response. The models were constructed using the lmer function in the lme4 package for R (Bates *et al.* 2015). Model selection was performed using the information-theoretic approach (AIC) proposed by Burnham and Anderson (2002). All possible combinations of the global model were analysed using the dredge function in the MuMIn package for R (Bartoń 2016). Only the models with  $\Delta AIC \leq 2$  are discussed, because they are treated as being equally supported (Burnham & Anderson 2002). Multiple competing models were assessed with regard to their fit to the data using AIC as the leading criterion, and the models with the lowest AIC were selected as the best fitting models. All data were analysed in the R environment (R Core Team 2019). The values reported are the mean  $\pm 1$  SE. Only those results with a probability of  $\alpha \leq 0.05$  are assumed to be statistically significant.

### 3. Results

During the ten years of the study, we found from 1 to 7 fledglings in the nest boxes (mean = 4.5, *SD* = 1.15, *N* = 319). The FED period ranged from 21 days in 2013 to 58 in 2019. Five models based on Akaike's information criterion achieved a value of  $\Delta AIC \leq 2$ . These models contained seven predictors, but three of these were the same in all the models (Table 2). The first model thus included

the FED, the mean and maximum temperatures and was selected as the best model. The four other weather parameters, i.e. total precipitation, number of days with precipitation, wind speed and minimum temperature, and also the habitat and location categories were excluded from the subsequent analysis.

The GLMM model showed that all three selected factors affected the reproduction of Common Kestrel (Table 3). Both the first egg date and the maximum temperature negatively influenced the number of fledglings. On the other hand, the mean daily temperature had a positive effect on Common Kestrel reproduction (Table 3).

### 4. Discussion

Our study showed that the first egg laying date and two weather factors but not the location or habitat type affected the reproduction of Common Kestrel. The first egg laying date tends to be negatively correlated with fledging success in birds: as the breeding season advances, environmental conditions deteriorate (Tolonen & Korpimäki 1995, Golawski & Meissner 2008). However, in a Mediterranean population of Common Kestrel, the timing of breeding was correlated with the amount of rainfall and temperature during the laying period (Costantini *et al.* 2010a). This relationship was explained by the birds exhibiting a certain plasticity, enabling them to adapt their laying time to environmental conditions: this was manifested by delayed egg-laying in rainier and colder breeding seasons. Variation in the mean laying date among breeding seasons suggests

Table 3. Estimated model coefficients for the best GLMM model of factors affecting the numbers of Common Kestrel *Falco tinnunculus* fledglings in eastern Poland.

Fixed effects	Estimate	Standard Error	<i>t</i> value	<i>P</i> value
Intercept	6.771	0.981	6.902	<0.001
First egg laying date	-0.023	0.005	-5.006	<0.001
Tmean	0.146	0.058	2.513	0.013
Tmax	-0.143	0.046	-3.090	0.002
Random effects	Variance	Standard deviation		
Nest box	0.043	0.206		
Residual	1.123	1.060		

that this species is sensitive to weather conditions (Carrillo & González-Dávila 2010a). Studies of different Common Kestrel populations across the Western Palearctic showed that it bred earlier where winter and spring temperatures were higher (Carrillo & González-Dávila 2010b).

Despite our assumption that trees should offer a superior microclimate in the canopy, nest boxes placed on utility poles had the same numbers of fledglings. The microclimate at breeding sites has been found to be a crucial factor for breeding success in many birds (e.g. Dawson *et al.* 2005; Ardia *et al.* 2006). Thus, microclimatic conditions during offspring development can variously affect hatchling mass, nestling growth, body condition and immunity; in extreme cases, heat stress and dehydration can lead to mortality (Murphy 1985; Bull 2003). However, we found a negative effect of maximum temperature on the number of fledglings, possibly because they are sensitive to high temperatures, which also may reduce food availability due, for example, to drought. This weather factor appeared to affect the broods in both locations to a similar extent; the tree canopy did not significantly improve conditions. Relationships between temperature and breeding parameters were found in American Kestrel: hatching success was lower in south-facing boxes, which were warmer than west-facing ones (Butler *et al.* 2009). However, a Finnish study of Common Kestrel did not corroborate these differences in the number of fledglings with respect to orientation (Valkama & Korpimäki 1999). It should be emphasised that in the cooler climate of Finland, high temperatures may have a different effect on

Common Kestrel reproduction than in Poland. Be that as it may, air temperatures in recent years have been rising quite distinctly (Ballester *et al.* 2010), so their effect on falcon nestlings 20 years ago may well have been different to what is happening now. Mean daily temperature seems to be conducive to greater breeding success in Common Kestrel: this could be related to the availability of prey, especially insects. *Orthopterans* are a very important component of the Common Kestrel diet (Gropalli 1992) as their development and activity are strongly dependent on temperature (Lactin & Johnson 1998, Maeno & Tanaka 2010). On the other hand, the costs of thermoregulation in cold weather may also increase food demands and affect reproductive parameters (see Carrillo & González-Dávila 2005). In general, therefore, higher temperatures may be very beneficial for falcon reproduction (Jenkins 2000, Rodríguez & Bustamante 2003).

In the present study, rainfall and wind were not important weather parameters. However, in most falcon species studied, precipitation does have a negative influence (Dawson & Bortolotti 2000, Charter *et al.* 2007b, Kreiderits *et al.* 2016). Such an adverse effect of rainfall on the number of fledglings has also been reported in shrikes *Laniidae*, whose diet is similar to that of Common Kestrel (Golawski 2006, Collister & Wilson 2007). Precipitation reduces the activity of the potential victims of birds, especially insects (Yosef 2000). A positive relationship between the total annual rainfall and Lesser Kestrel reproduction was found in Spain, which was expected, because rainfall is the main factor



limiting productivity in Mediterranean climates (Rodríguez & Bustamante 2003). A warmer and rainier spring also enhanced fledgling success in an Italian population of Common Kestrel (Costantini *et al.* 2010a). Geographically related conditions could be the main explanation for the differences in breeding effort here (Carrillo & González-Dávila 2010b).

Nest boxes dedicated to kestrels and deployed on trees are often taken over by other bird species like owls (Golawski *et al.* unpublished data). Moreover, this type of location is exposed to higher predation pressure on the part of Pine Martens (Valkama & Korpimäki 1999). Utility poles therefore appear to be a safer location than trees. Assuming further that both locations had the same reproductive parameters and that nest boxes on utility poles are easier to monitor and maintain, the latter should be the preferred form of location in conservation programmes. Moreover, where habitat types form a large mosaic, as in eastern Poland, the surface area of the main habitat type will not be a decisive factor affecting kestrel productivity. Most of the feeding grounds will probably contain the necessary environmental components ensuring food abundance (see Golawski & Meissner 2008).

In conclusion, our study showed that the number of Common Kestrel fledglings in nest boxes attached to utility poles was the same as in those deployed on trees. Also, there was no clear link with the nest box location, weather factors and the dominant habitat type. These findings may be very useful in ongoing conservation programmes because nest boxes on utility poles are easier to monitor and maintain and may be safer than nest boxes on trees.

### *Sääolosuhteiden ja pesäpöntön sijainnin vaikutus tuulihaukan lisääntymiseen Puolassa*

Keinopesiä, kuten pesäpönttöjä, käytetään usein parantamaan lintujen lisääntymismenestystä, ja ne mahdollistavat lajin ekologian ja evoluution tutkimisen. Tutkimme, miten pesäpönttöjen sijainti ja paikalliset sääolosuhteet vaikuttavat tuulihaukan lisääntymismenestykseen (poikastuotto) Puolassa. Tutkimuksen otoskoko oli 173 pesäpönttöä ja 319 poikuetta. Analysoimme kuutta

meteorologista parametriä, joista keskimääräinen ja maksimilämpötila valikoituivat parhaaseen tilastomalliin: päivän keskilämpötilalla oli positiivinen vaikutus tuulihaukan pesintään, kun taas maksimilämpötilan vaikutus oli päinvastainen. Muninnan aloitus korreloi positiivisesti poikastuoton kanssa. Oletimme, että puiden latvustossa sijaitsevilla pöntöillä mikroilmasto olisi suotuisampi pesinnän kannalta kuin sähköpylväeseen kiinnitetyssä pöntössä, mutta pesimämenestys ei eronnut näissä kahdessa ryhmässä. Tulokset auttavat lajin suojelussa, ja muun muassa varmistavat että lajin populaatioseurannassa voidaan suosia sähköpylväisiin kiinnitettyjä keinopesiä, jotka helpottavat seurantaa.

*Acknowledgements.* We would like to thank all the volunteers for their dedicated help in the field work, in particular, Przemysław Abramowski, Andrzej Borowski, Maciej Cmoch, Przemysław Kurdej, Janusz Łuczak, Andrzej Łukijańczuk, Jerzy Mikołajczuk, Przemysław Obłóza, Wojciech Okliński, Agnieszka Parapura, Ryszard Ryś, Monika Stefaniak, Marcin Stępień, Piotr Szweczyk, Adam Tarłowski and Grażyna Zabłocka.

## References

- Altwegg, R., Jenkins, A. & Abadi, F. 2014: Nestboxes and immigration drive the growth of an urban Peregrine Falcon *Falco peregrinus* population. *Ibis* — 156: 107–115.
- Ardia, D. R., Perez, J. H., & Clotfelter, E. D. 2006: Nest box orientation affects internal temperature and nest site selection by Tree Swallows. — *Journal of Field Ornithology* 77: 339–344.
- Ballester, J., Rodó, X., & Giorgi, F. 2010: Future changes in Central Europe heat waves expected to mostly follow summer mean warming. — *Climate Dynamics* 35: 1191–1205.
- Bartoń, K. 2016: MuMIn: Multi-Model Inference. — R package version 1.15.6: <https://cran.r-project.org/web/packages/MuMIn>.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015: Fitting linear mixed-effects models using lme4. — *Journal of Statistical Software* 67: 1–48.
- Bortolotti, G. R. 1994: Effects of nest-box size on nest-site preference and reproduction in American kestrels. — *Journal of Raptor Research* 28: 127–133.
- Bull, E. L. 2003: Use of nest boxes by Vaux's Swifts. — *Journal of Field Ornithology* 74: 394–400.
- Burnham, K.P., Anderson, D.R. 2002: Model selection and multimodel inference. — Heidelberg: Springer.
- Butler, M.W., Whitman, B. A., & Duffy, A. M. Jr. 2009: Nest box temperature and hatching success of American Kes-

- trels varies with nest box orientation. — *Wilson Journal of Ornithology* 121: 778–782.
- Bux, M., Giglio, G. & Gustin, M. 2008: Nest box provision for Lesser Kestrel *Falco naumanni* populations in the Apulia region of southern Italy. — *Conservation Evidence* 5: 58–61.
- Carlzon, L., Karlsson, A., Falk, K., Liess, A. & Møller, S. 2018: Extreme weather affects Peregrine Falcon (*Falco peregrinus tundrius*) breeding success in South Greenland. — *Ornis Hungarica* 26: 38–50.
- Carrillo, J. & González-Dávila, E. 2005: Breeding biology and nest characteristics of the Eurasian Kestrel in different environments on an Atlantic island. — *Ornis Fennica* 82: 55–62.
- Carrillo, J. & González-Dávila, E. 2010a: Impact of weather on breeding success of the Eurasian Kestrel *Falco tinnunculus* in a semi-arid island habitat. — *Ardea* 98: 51–58.
- Carrillo, J. & González-Dávila, E. 2010b: Geo-environmental influences on breeding parameters of the Eurasian Kestrel (*Falco tinnunculus*) in the Western Palearctic. — *Ornis Fennica* 87: 15–25.
- Carstens, K. F., Kassanjee, R., Little, R. M., Ryan, P. G., & Hockey, P. A. R. 2019: Breeding success and population growth of Southern Ground Hornbills *Bucorvus leadbeateri* in an area supplemented with nest boxes. — *Bird Conservation International*, <https://doi.org/10.1017/S0959270919000108>
- Charter, M., Izhaki, I., Bouskila, A., Leshem, Y. & Penteriani, V. 2007a: Breeding success of the Eurasian Kestrel (*Falco tinnunculus*) nesting on buildings in Israel. — *J. Raptor Res.* 41: 139–143.
- Charter, M., Izhaki, I., Bouskila, A., and Leshem, Y. 2007b: The effect of different nest types on the breeding success of Eurasian Kestrels (*Falco tinnunculus*) in a rural ecosystem. — *Journal of Raptor Research* 41: 143–149.
- Collister, D. M. & Wilson, S. 2007: Contributions of weather and predation to reduced breeding success in a threatened northern loggerhead shrike population. — *Avian Conservation and Ecology* 2: article 11.
- Costantini, D., Carello, L., & Dell’Omo, G. 2010a: Patterns of covariation among weather conditions, winter North Atlantic Oscillation index and reproductive traits in Mediterranean kestrels. — *Journal of Zoology* 280: 177–184.
- Costantini, D., Carello, L. & Dell’Omo, G. 2010b: Temporal covariation of egg volume and breeding conditions in the common kestrel (*Falco tinnunculus*) in the Mediterranean region. — *Ornis Fennica* 87: 144–152.
- Costantini, D. & Dell’Omo, G. 2020: The kestrel: ecology, behaviour and conservation of an open-land predator. Cambridge University Press, Cambridge.
- Dawson, R. D., Lawrie, C. C., & O’Brien, E. L. 2005: The importance of microclimate variation in determining size, growth and survival of avian offspring: experimental evidence from a cavity nesting passerine. — *Oecologia* 144: 499–507.
- Dawson, R. D. & Bortolotti, G.R. 2000: Reproductive success of American kestrels: the role of prey abundance and weather. — *Condor* 102: 814–822.
- Degirmendzic, J., Kozuchowski, K. & Zmudzka, E. 2004: Changes of air temperature and precipitation in Poland in the period 1951–2000 and their relationship to atmospheric circulation. — *International Journal of Climatology* 24: 291–310.
- Fargallo, J. A., Blanco, G., Potti, J. & Viñuela, J. 2001: Nest-box provisioning in a rural population of Eurasian Kestrels: breeding performance, nest predation and parasitism. — *Bird Study* 48: 236–244.
- Golawski, A. 2006: Impact of weather on partial loss of nestlings in the Red-backed Shrike *Lanius collurio* in eastern Poland. — *Acta Ornithologica* 41: 15–20.
- Golawski, A., Kasprzykowski, Z. & Kowalski, M. 2003: The occurrence of the Barn Owl *Tyto alba* in sacred buildings in central-eastern Poland. — *Ornis Hungarica* 12-13: 275–277.
- Golawski, A. & Meissner, W. 2008: The influence of territory characteristics and food supply on the breeding performance of the Red-backed Shrike (*Lanius collurio*) in an extensively farmed region of eastern Poland. — *Ecological Research* 23: 347–353.
- Gropalli, R. 1992: Sull’ alimentazione insettivora del Gheppio *Falco tinnunculus* e del Falco cuculo *Falco vesperinus*. — *Picus* 18: 29–31. (In Italian with English summary)
- Jenkins, A. R. 2000: Factors affecting breeding success of Peregrine and Lesser Falcons in South Africa. — *Ostrich* 71: 385–392.
- Klein, A., Nagy, T., Csorgo, T. & Matics, R. 2007: Exterior-nest-boxes may negatively affect Barn Owl *Tyto alba* survival: an ecological trap. — *Bird Conservation International* 17: 273–281.
- Korpimäki E., Lahti K., May C.A., Parkin D. T., Powell G.B., Tolonen P., Wetton J.H. 1996: Copulatory behavior and paternity determination by DNA fingerprinting in Kestrel: effects of cyclic food abundance. — *Animal Behaviour* 51: 945–955.
- Kreiderits, A., Gamauf, A., Krenn, H. & Sumasgutner, P. 2016: Investigating the influence of local weather conditions and alternative prey composition on the breeding performance of urban Eurasian Kestrels (*Falco tinnunculus*). — *Bird Study* 63: 369–379.
- Lactin, D. J. & Johnson, D. L. 1998: Environmental, physical, and behavioural determinants of body temperature in grasshopper nymphs (Orthoptera: Acrididae). — *Canadian Entomologist* 130: 551–577.
- Lambrechts M.M., Wiebe K.L., Sunde P., Solonen T., Sergio F., Roulin A., Møller A.P., López B.C., Fargallo J.A., Exo M., Dell’Omo G., Costantini D., Charter M., Butler M.W., Bortolotti G.R., Arlettaz R. & Korpimäki E., 2012. Nest-box design for the study of diurnal raptors and owls is still an overlooked point in ecological, evolutionary and conservation studies: a review. — *Journal of Ornithology* 153, 23–34.

- Le Roux, D. S., Ikin, K., Lindenmayer, D. B., Bistricher, G., Manning, A. D. & Gibbons, P. 2016: Effects of entrance size, tree size and landscape context on nest box occupancy: Considerations for management and biodiversity offsets. — *Forest Ecology and Management* 366: 135–142.
- Libois, E., Gimenez, O., Oro, D., Minguéz, E., Pradel, R. & Sanz-Aguilar, A. 2012: Nestboxes: a successful management tool for the conservation of an endangered seabird. — *Biological Conservation* 155: 39–43.
- Maeno K., & Tanaka S. 2010: Patterns of nymphal development in the desert locust, *Schistocerca gregaria*, with special references to phase-dependent growth and extra molting. — *Applied Entomology and Zoology* 45: 513–519.
- Mainwaring, M. C. 2015: The use of man-made structures as nesting sites by birds: A review of the costs and benefits. — *Journal for Nature Conservation* 25: 17–22.
- Martínez-de la Puente, J., Merino, S., Lobato, E., Aguilar, J. R., Cerro, S., Castañeda, R. R., & Moreno, J. 2009: Does weather affect biting fly abundance in avian nests? — *Journal of Avian Biology* 40: 653–657.
- McDonald, P. G., Olsen, P. D. & Cockburn A. 2004: Weather dictates reproductive success and survival in the Australian Brown Falcon *Falco berigora*. — *Journal of Animal Ecology* 73: 683–692.
- Murphy, M. T. 1985: Nestling Eastern Kingbird growth: effects of initial size and ambient temperature. — *Ecology* 66: 162–170.
- Norrdahl K., Korpimäki E. 2002. Seasonal changes in the numerical response of predators to cyclic vole populations. — *Ecography* 25: 428–438.
- R Core Team. 2019: R: a language and environment for statistical computing. Vienna: The R Foundation for Statistical Computing. — <https://www.R-project.org/>.
- Rodríguez, J., Avilés, J. M. & Parejo, D. 2011: The value of nestboxes in the conservation of Eurasian rollers *Coracias garrulus* in southern Spain. — *Ibis* 153: 735–745.
- Rodríguez, C. & Bustamante, J. 2003: The effect of weather on lesser kestrel breeding success: can climate change explain historical population declines? — *Journal of Animal Ecology* 72: 793–810.
- Rohrbaugh, R. W. Jr., & Yahner, R. H. 1997: Effects of macrohabitat and microhabitat on nest-box use and nesting success of American Kestrels. — *Wilson Bulletin* 109: 410–423.
- Schaub, T., Meffert, P. J., & Kerth, G. 2016: Nest-boxes for common swift *Apus apus* as compensatory measures in the context of building renovation: Efficacy and predictors of occupancy. — *Bird Conservation International* 26: 164–176.
- Smallwood, I. A. & Collopy, M. W. 2009: Southeastern American Kestrels respond to an increase in the availability of nest cavities in north-central Florida. — *Journal of Raptor Research* 43: 291–300.
- Smallwood, J. A., Causey, M. F., Mossop, D. H., Klucsarits, J. R., Robertson, B., Robertson, S., Mason, J., Maurer, M. J., Melvin, R. J., Dawson, R. D., Bortolotti, G. R., Parrish, J. W., Breen, T. F. & Boyd, K. 2009: Why are American kestrel (*Falco sparverius*) populations declining in North America? Evidence from nest-box programs. — *Journal of Raptor Research* 43: 274–282.
- Statistical Yearbook 2015: Mazowieckie Voivodship 2015. Warszawa: Statistical Office in Warszawa.
- Sumasgutner, P., Jenkins, A., Amar, A. & Altwegg, R. 2020: Nest boxes buffer the effects of climate on breeding performance in an African urban raptor. — *PloS ONE* 15(6): e0234503.
- Tolonen, P., & Korpimäki, E. 1995: Parental effort of kestrels (*Falco tinnunculus*) in nest defence: effects of laying time, brood size, and varying survival prospects of offspring. — *Behavioural Ecology* 6: 435–441.
- Valkama, J. & Korpimäki, E. 1999: Nestbox characteristics, habitat quality and reproductive success of Eurasian Kestrels. — *Bird Study* 46: 81–88.
- Yosef, R. 2000: Nesting ecology of the Loggerhead Shrike (*Lanius ludovicianus*) in southcentral Florida. — *Ring* 22: 111–121.